



Handläggande organ

Fastställt

Utgåva

Sida

Standardiseringsgruppen STG

1999-12-22

1

1 (1+19)

© Copyright SIS. Reproduction in any form without permission is prohibited.

## Soil quality – Inhibition of reproduction of *Collembola* (*Folsomia candida*) by soil pollutants

This Swedish standard consists of the English version of ISO 11267:1999, Soil quality – Inhibition of reproduction of *Collembola* (*Folsomia candida*) by soil pollutants.

This International Standard describes a method for determining the effects of substances on the reproduction of *Folsomia candida* by dermal and alimentary uptake in a defined artificial soil substrate.

This method is not applicable to volatile substances.

Swedish Standards corresponding to documents referred to in this Standard are listed in "Catalogue of Swedish Standards", issued by SIS.

## Markundersökningar – Reproduktionshämning av *Collembola* (*Folsomia candida*) på grund av markföroreningar

Denna standard utgörs av den engelska versionen av den internationella standarden ISO 11267:1999, Soil quality – Inhibition of reproduction of *Collembola* (*Folsomia candida*) by soil pollutants.

Denna standard beskriver en metod för att bestämma substansers reproduktionshämmande effekt på *Folsomia candida* genom hud- och näringsupptag från en definierad artificiell jord.

Metoden är inte användbar för flyktiga substanser.

I SIS "Katalog över svensk standard" framgår vilka av de publikationer som omnämns i denna standard som har fastställts som svensk standard och som har översatts till svenska.

ICS 13.080.00

**ISO 11267:1999(E)**

**Contents**

**1 Scope ..... 1**

**2 Normative references ..... 1**

**3 Terms and definitions ..... 1**

**4 Principle ..... 2**

**5 Reagents ..... 2**

**6 Apparatus ..... 3**

**7 Procedure ..... 3**

**8 Calculation and expression of results ..... 5**

**9 Validity of the test ..... 6**

**10 Test report ..... 7**

**Annex A (informative) Techniques for rearing and breeding of Collembola ..... 8**

**Annex B (informative) Techniques for counting juvenile springtails ..... 11**

**Annex C (informative) Determination of water-holding capacity of artificial soil ..... 12**

**Annex D (informative) Guidance on adjustment of pH of artificial soil ..... 13**

**Annex E (informative) Determination of effects of contaminated soil on Collembola reproduction ..... 14**

**Bibliography ..... 16**

© ISO 1999

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the publisher.

International Organization for Standardization  
 Case postale 56 • CH-1211 Genève 20 • Switzerland  
 Internet iso@iso.ch

Printed in Switzerland

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

International Standard ISO 11267 was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

Annexes A to E of this International Standard are for information only.

## Introduction

This International Standard describes a method for testing the effects of chemicals on the reproduction of Collembola in artificial soil. It can be adapted for use for testing or comparing soils to assess, for example, the effects of remediation treatments, and for assessing sublethal effects and no-effect levels for pesticides or other added chemicals.

# Soil quality — Inhibition of reproduction of *Collembola* (*Folsomia candida*) by soil pollutants

## 1 Scope

This International Standard describes a method for determining the effects of substances on the reproduction of *Folsomia candida* by dermal and alimentary uptake in a defined artificial soil substrate.

The method is not applicable to volatile substances, i.e. substances for which H (Henry's constant) or the air/water partition coefficient is greater than 1, or for which the vapour pressure exceeds 0,013 3 Pa at 25 °C.

NOTE 1 The stability of the test substance cannot be assured over the test period. No allowance is made in the test method described for possible degradation of the test substance over the course of the experiment..

NOTE 2 Recommendations for adapting the method for comparing or monitoring soil quality are given in annex E.

## 2 Normative references

The following normative documents contain provisions which, through reference in this text, constitute provisions of this International Standard. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the normative documents indicated below. For undated references, the latest edition of the normative document referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO 10390:1994, *Soil quality — Determination of pH*.

ISO 11268-1:1993, *Soil quality — Effects of pollutants on earthworms (Eisenia fetida) — Part 1: Determination of acute toxicity using artificial soil substrate*.

ISO 11274, *Soil quality — Determination of the water retention characteristic — Laboratory methods*.

## 3 Terms and definitions

For the purposes of this International Standard, the following terms and definitions apply.

### 3.1

#### LOEC

#### lowest observed effect concentration

lowest concentration of the test substance which is observed to have a significant effect when compared with the control

NOTE All test concentrations above the LOEC have a harmful effect equal to, or greater than that observed at the LOEC.

**3.2**

**NOEC**

**no observed effect concentration**

highest tested concentration of a test substance at which no lethal or other effect (such as mass alteration) is observed

**3.3**

**EC10**

concentration estimated to reduce the reproduction rate at the end of the test by 10 % compared to the control

NOTE All effect concentrations are expressed as mass of test substance per dry mass of the test substrate (5.2.1).

**3.4**

**EC50**

concentration estimated to reduce the reproduction rate at the end of the test by 50 % compared to the control

NOTE All effect concentrations are expressed as mass of test substance per dry mass of the test substrate (5.2).

**3.5**

**reproduction**

increase in the mean numbers of offspring in each test vessel after 28 days incubation under the specified test conditions

**4 Principle**

The effects of different concentrations of test substance on the reproduction of 10- to 12-day old springtails (*Folsomia candida*) in a defined artificial substrate are determined. The springtails are incubated until offspring ( $F_1$ ) emerge from eggs laid by mature adults, and the number of offspring are determined. Normally offspring emerge after 28 days in control experiments.

**5 Reagents**

**5.1 Biological material.**

In this test, 10- to 12-day old juvenile springtails of the species *Folsomia candida* (Willem) are used (see A.1. for details on synchronization of breeding).

**5.2 Test substrate**, consisting of the wet basic soil substrate (defined artificial soil in accordance with ISO 11268-1) the test substance and deionized water.

**5.2.1 Soil substrate.**

Component	Dry mass fraction
Sphagnum peat, air-dried, finely ground and with no visible plant remains	10 %
Kaolinite clay (air-dry) containing not less than 30 % kaolinite	20 %
Industrial quartz sand (air dried, and predominantly fine sand with more than 50 % of particle size 0,05 mm to 0,2 mm)	70 %

Add sufficient (usually 0,5 % to 1 %) calcium carbonate ( $\text{CaCO}_3$ ), pulverized and of recognized analytical grade, to bring the pH (as measured in 1 mol/l KCl solution) to  $6,0 \pm 0,5$  at the start of the test. The pH shall be determined in accordance with ISO 10390.

NOTE 1 The amount of calcium carbonate required will depend on the components of the soil substrate and should be determined by measurements on subsamples (see annex D) immediately before the test.

The dry constituents are blended in the correct proportions and mixed thoroughly in a large-scale laboratory mixer or a household mixer. A portion of the deionized water required is added while mixing is continued. This is the basic soil substrate. The overall water content of the test and control substrates respectively should be adjusted to give the substrate a crumbly structure to enable springtails to penetrate substrate cavities. This is normally at a water content of 40 % to 60 % of the total water-holding capacity determined in accordance with ISO 11274, or with the method given in annex C. Water content and pH are determined before the start of the test and at the end of the test for the control and for each concentration tested.

NOTE 2 Allowance should be made for any water or quartz sand used for introducing the test substance into the soil.

**5.2.2 Control substrate**, consisting of the basic substrate and deionized water. If the preparation of the test requires the use of an auxiliary agent, then an additional control will be necessary.

### 5.3 Reference substance.

To ensure the quality of the test system, tests should be performed regularly (once or twice a year) with a reference substance.

The agricultural chemicals Betanal plus (a.i. 160 g/l Phenmedipham) and E 605 forte (a.i. 507,5 g/l Parathion) have been tested in a ring test, and are recommended as reference substances.

**WARNING — When handling these chemicals, appropriate precautions should be taken to avoid ingestion or skin contact.**

NOTE Betanal plus: Effects on reproduction ( $\alpha = 0,05$ ) were observed at concentrations of between 100 mg and 200 mg of the product per kilogram dry mass of the substrate.

E 605 forte: Effects on mortality and reproduction were observed at concentrations of between 0,18 mg and 0,32 mg product and between 0,1 mg and 0,18 mg product per kilogram dry mass of the substrate respectively.

## 6 Apparatus

Standard laboratory equipment, and:

**6.1 Glass containers** (able to be closed tightly) of about 100 ml capacity and with a diameter of about 5 cm.

**6.2 Apparatus** capable of measuring pH and water content of the substrate.

**6.3 Exhaustor** for transfer of springtails (see A.2).

**6.4 Enclosure**, capable of being controlled to a temperature of  $20\text{ °C} \pm 2\text{ °C}$ .

**6.5 Light source**, capable of delivering a constant light intensity of 400 lx to 800 lx at the substrate surface at a controlled light:dark cycle of between 12 h:12 h and 16 h: 8 h.

## 7 Procedure

### 7.1 Preparation of the test

#### 7.1.1 Test concentrations

a) NOEC approach:

At least five concentrations in a geometric series at a factor not exceeding 2 (e.g.  $\sqrt[4]{10} \sim 1,8$ ) should be selected to give an estimation of the LOEC/NOEC of the reproduction rate.

b)  $EC_x$  approach:

For the  $EC_x$  approach, a higher number of concentrations should be used (e.g. 12), and each tested with two replicates. The spacing of concentrations may be variable (i.e. smaller at low concentrations, and larger at high concentrations).

The concentrations of the test substance shall be expressed as mass of substance per dry mass of soil substrate (mg/kg) (see 5.2).

If no other relevant toxicity data are available, the concentrations selected to provide the LOEC/NOEC or  $EC_{10}$  will be based on the results of a preliminary test (see 7.2).

## 7.1.2 Introduction of the test substance

### 7.1.2.1 Water-soluble substances

Immediately before starting the test, dissolve the quantity of test substance in the water required for the replicates of one concentration (or that portion of it necessary to wet the soil substrate in order to meet the requirements of 5.2.1), and mix it thoroughly with the basic soil substrate before introducing it into the test containers.

Continue as described in 7.1.3.

### 7.1.2.2 Substances insoluble in water but soluble in organic solvents

Dissolve the quantity of test substance required to obtain the desired concentration in a volatile solvent (such as acetone or hexane) and mix it with a portion of the quartz sand required. After having evaporated the solvent by placing the container in a fume hood, add the remainder of the basic substrate (allowing for the amount of sand used to prepare the test substance) and the water, and mix thoroughly before introducing it into the test containers.

Continue as described in 7.1.3.

**NOTE** Ultrasonic dispersion, organic solvents, emulsifiers or dispersants may be used to disperse substances with low aqueous solubility. When such auxiliary substances are used, all test concentrations and an additional control should contain the same minimum amount of auxiliary substance.

**WARNING** — Appropriate precautions should be taken when dealing with solvent vapour to avoid danger from inhalation or explosion, and to avoid damage to extraction equipment, pumps etc.

### 7.1.2.3 Substances insoluble in water or organic solvents

For a substance insoluble in a volatile solvent, prepare a mixture of 10 g of finely ground industrial quartz sand (see 5.2) and the quantity of the test substance required to obtain the desired concentration. Add the remainder of the basic substrate (allowing for the amount of sand used to prepare the test substance) and the water, and mix it thoroughly before introducing it into the test containers.

Continue as described in 7.1.3.

## 7.1.3 Introduction of the test organisms

Ten juvenile springtails (10 to 12 days old) are placed in each test container.

Springtails are tapped or sucked from the breeding containers to transfer them to the test containers. This can easily be done using an exhaustor as described in clause A.2. Before they are transferred to the test containers, organisms are counted and checked for damage both to reduce control mortality and to avoid systematic trial errors.

### 7.1.4 Control container

Prepare control containers in the same way as test containers, but without addition of the test substance. If the preparation of the test requires the use of auxiliary substances (see 7.1.2.2), use additional control containers. Treat these containers in the same way as those without the test substance.